

Supplementary material

Spatial patterns of wildfire ignitions in south-eastern Australia

Kathryn M. Collins^{A,C}, Owen F. Price^A and Trent D. Penman^{A,B}

^ACentre for Environmental Risk Management of Bushfires, University of Wollongong, Wollongong, NSW 2522, Australia.

^BSchool of Ecosystem and Forest Science, University of Melbourne, Water Street, Creswick, Vic. 3363, Australia.

^CCorresponding author. Email: kmc624@uowmail.edu.au




Fig. S1. Major agro-climatic zones in the study area in relation to subregions (source: Hutchinson *et al.* 2005).




Fig. S2. Predominant natural vegetation in relation to subregions. (Source: National Vegetation Information System version 4.1, Department of the Environment, available at <http://www.environment.gov.au/land/native-vegetation/national-vegetation-information-system/data-products>, accessed 28 August 2015.)




Fig. S3. Land-use classification classes in relation to subregions. Intensive land use includes residential and industrial areas. (Source: The Land Use of Australia, Version 4, 2005–06, Australian Bureau of Agricultural and Resource Economics and Sciences (ABARES), available at <http://www.agriculture.gov.au/abares/aclump/Pages/land-use/data-download.aspx>, accessed 28 August 2015.)