

Supplementary Material

Current insecticide treatments used in locust control have less of a short-term impact on Australian arid-zone reptile communities than does temporal variation

Kimberly Maute^{A,E}, Kristine French^A, C. Michael Bull^B, Paul Story^{A,C} and Grant Hose^D

^ASchool of Biological Sciences, University of Wollongong, Wollongong, NSW 2522, Australia.

^BSchool of Biological Sciences, Flinders University, Adelaide, SA, Australia.

^CAustralian Plague Locust Commission, Fyshwick, ACT, Australia.

^DSchool of Biological Sciences, Macquarie University, Sydney, NSW, Australia.

^ECorresponding author. Email: kmaute@uow.edu.au

Table S1: List of reptile species and numbers of captures during the two year study (2012 – 2014) at Fowlers Gap Arid Zone Research Station, NSW Australia. Reptile species are listed in the order of abundance.

Reptile Species	Number of captures
<i>Ctenotus strauchii</i>	118
<i>Ctenotus leonhardii</i>	50
<i>Tympanocryptis tetraporophora</i>	28
<i>Ctenotus schomburgki</i>	14
<i>Ctenotus olympicus</i>	12
<i>Menetia greyii</i>	12
<i>Heteronotia binoei</i>	9
<i>Rhynchoedura spp</i>	8
<i>Diplodactylus tessellatus</i>	6
<i>Pogona vitticeps</i>	5
<i>Morethia adelaidensis</i>	5
<i>Delma tincta</i>	4
<i>Ctenotus regius</i>	4
<i>Suta suta</i>	3
<i>Eremiascincus fasciolatus</i>	2
<i>Morethia boulergeri</i>	2
<i>Varanus gouldii</i>	2
<i>Ctenophorus nuchalis</i>	1
<i>Gehyra variegata</i>	1
<i>Lucasium byrnie</i>	1
<i>Nephurus levis</i>	1
<i>Pseudonaja textilis</i>	1