

10.1071/WR16065_AC

©CSIRO 2016

Supplementary Material: *Wildlife Research* **43**, 590–598

SUPPLEMENTARY MATERIAL

Creeping into a wild boar stomach to find traces of supplementary feeding

Ježek Miloš^A, Holá Michaela^{A,B}, Kušta Tomáš^A and Červený Jaroslav^A

^ADepartment of Game Management and Wildlife Biology, Faculty of Forestry and Wood Sciences, Czech University of Life Sciences Prague, Kamýcká 129, Prague, 165 21, Czech Republic.

^BCorresponding author: Email: mhola@fld.czu.cz

Table S1. Frequency of occurrence (%) of all diet items identified in wild boar stomachs collected from hunted individuals in four study sites in the Czech Republic during 2010–2012

DIET ITEM	LATIN NAME	SPRING	SUMMER	WINTER
Corn	<i>Zea mays</i>	35%	26%	48%
Oat	<i>Avena sativa</i>	1%	0%	12%
Wheat	<i>Triticum aestivum</i>	21%	41%	30%
Barley	<i>Hordeum vulgare</i>	15%	2%	15%
Cereals unspecified		4%	31%	17%
CEREALS TOTAL		55%	59%	76%
Rape	<i>Brassica napus</i>	4%	2%	3%
Sunflower	<i>Helianthus annuus</i>	3%	0%	5%
Potatos	<i>Solanum</i>	0%	2%	0%
Beet	<i>Beta vulgaris</i>	0%	0%	3%
OILSEEDS AND ROOT CROPS TOTAL		7%	2%	8%
Fresh grass		46%	50%	30%
Hay		0%	2%	6%
GRASSES TOTAL		46%	52%	36%
Nettle	<i>Urtica dioica</i>	1%	0%	1%
Dandelion	<i>Taraxacum officinale</i>	18%	7%	2%
Plantain	<i>Plantago lanceolata</i>	3%	2%	1%
Herbs unspecified		4%	4%	1%
HERBS TOTAL		23%	11%	5%
Roots		39%	41%	45%
Rhizomes		11%	6%	6%
ROOT AND RHIZOMES TOTAL		44%	43%	47%
Leaves of oak	<i>Quercus robur</i>	0%	4%	7%
Leaves of beech	<i>Fagus sylvatica</i>	0%	4%	1%
Leaves of aspen	<i>Populus tremula</i>	0%	4%	1%
Leaves of maple	<i>Acer platanoides</i>	1%	2%	0%
Leaves of birch	<i>Betula</i>	0%	0%	2%
Leaves of alder	<i>Alnus</i>	0%	2%	0%
Leaves of hornbeam	<i>Carpinus</i>	1%	0%	0%
LEAVES OF TREES TOTAL		3%	15%	9%

Table S1_Part 2

DIET ITEM	LATIN NAME	SPRING	SUMMER	WINTER
Seeds of Fireweed	<i>Epilobium angustifolium</i>	1%	0%	0%
Seeds of Fallopia	<i>Fallopia</i>	7%	0%	6%
Seeds of Goosefoots	<i>Chenopodium</i>	3%	2%	2%
Seeds of Bindweed	<i>Convolvulus</i>	3%	9%	9%
Seeds of Bistorta	<i>Bistorta</i>	0%	0%	2%
Seeds of Common flax	<i>Linum usitatissimum</i>	1%	0%	0%
Seeds of Madder	<i>Galium</i>	8%	11%	7%
Seeds of Campion	<i>Silene</i>	1%	2%	0%
Seeds of Amaranth	<i>Amaranthus</i>	1%	2%	2%
Seeds of Sorrels	<i>Rumex</i>	0%	4%	2%
Seeds of Wood violet	<i>Viola</i>	1%	0%	1%
SEEDS OF HERBS TOTAL		14%	28%	15%
Seeds of Dog-rose	<i>Rosa canina</i>	3%	0%	1%
Seeds of Hawthorn	<i>Crataegus</i>	0%	0%	0%
Seeds of Sea buckthorns	<i>Hippophae</i>	0%	2%	0%
Seeds of Ash	<i>Fraxinus</i>	1%	0%	0%
Seeds of hornbeam	<i>Carpinus</i>	7%	0%	3%
Seeds of beech	<i>Fagus sylvatica</i>	1%	7%	6%
Seeds of oak	<i>Quercus robur</i>	25%	17%	17%
Seeds of Walnut trees	<i>Juglans</i>	0%	0%	0%
SEEDS OF TREES TOTAL		34%	22%	25%
Fruits of European blueberry	<i>Vaccinium myrtillus</i>	1%	7%	1%
Fruits of European raspberry	<i>Rubus idaeus</i>	3%	4%	0%
Fruits of Apple tree	<i>Malus</i>	4%	17%	16%
Fruits of Pear tree	<i>Pyrus</i>	0%	6%	1%
Fruits of Wild cherry	<i>Prunus avium</i>	3%	17%	1%
Fruits of Egg plum	<i>Prunus domestica</i>	1%	9%	0%
Fruits of Grape vine	<i>Vitis vinifera</i>	0%	0%	0%
FRUITS TOTAL		11%	44%	18%
Red deer	<i>Cervus elaphus</i>	0%	0%	0%
Roe deer	<i>Capreolus capreolus</i>	4%	2%	3%
Wild boar	<i>Sus scrofa</i>	0%	0%	2%
Wild hare	<i>Lepus europeus</i>	4%	0%	1%
Bank vole	<i>Clethrionomys glareolus</i>	1%	9%	1%
Yellow-necked mouse	<i>Apodemus flavicollis</i>	1%	0%	2%
Common vole	<i>Microtus arvalis</i>	1%	2%	3%
Field vole	<i>Microtus agrestis</i>	1%	0%	0%
Eurasian shrew	<i>Sorex araneus</i>	0%	2%	0%
European water vole	<i>Arvicola terrestris</i>	0%	2%	3%
MAMMALS TOTAL		13%	17%	14%

Table S1_Part 3

DIET ITEM	LATIN NAME	SPRING	SUMMER	WINTER
FISH TOTAL	<i>Osteichthyes</i>	0%	0%	3%
BIRDS TOTAL	<i>Aves</i>	6%	6%	2%
European common frog	<i>Rana temporaria</i>	1%	0%	0%
Slow worm	<i>Anguis fragilis</i>	1%	0%	0%
AMPHIBIANS TOTAL	<i>Amphibia</i>	3%	0%	0%
Ants	<i>Formicidae</i>	0%	2%	0%
Caterpillars		0%	0%	0%
Larvae		48%	19%	37%
Long-horned grasshoppers	<i>Ensifera</i>	0%	2%	0%
Reddish worm	<i>Lumbricus terrestris</i>	25%	17%	2%
Beetles	<i>Coleoptera</i>	28%	37%	6%
Spiders	<i>Araneae</i>	1%	0%	0%
Molluscs	<i>Mollusca</i>	14%	7%	4%
INVERTEBRATES TOTAL	<i>Invertebrata</i>	42%	50%	39%
MUSHROOMS TOTAL	Fungi	13%	4%	14%
OTHERS TOTAL		25%	15%	43%