

10.1071/CH13227\_AC

©CSIRO 2013

Australian Journal of Chemistry 2013, 66(10), 1253-1259

## Supplementary Material

### Nucleophilic aromatic substitution of bis(pentafluorophenyl)mercury with various bulky nucleophiles and the structures of [Hg(C<sub>6</sub>F<sub>4</sub>X-4)<sub>2</sub>] (X = *cyclo-C*<sub>5</sub>H<sub>10</sub>N, OCH(CH<sub>3</sub>)<sub>2</sub>, OC(CH<sub>3</sub>)<sub>3</sub>)

Glen B. Deacon <sup>A C</sup>, Peter C. Junk <sup>B C</sup> and Jenny Luu <sup>A</sup>


<sup>A</sup> School of Chemistry, Monash University Clayton, Victoria, 3800, Australia

<sup>B</sup> School of Pharmacy & Molecular Sciences, James Cook University, Townsville, Qld, 4811, Australia.


<sup>C</sup> Corresponding authors' email addresses: [Peter.Junk@jcu.edu.au](mailto:Peter.Junk@jcu.edu.au) [Glen.Deacon@monash.edu](mailto:Glen.Deacon@monash.edu)

**Figure S1:** **A:**  $^{19}\text{F}\{^1\text{H}\}$  NMR of the first attempts at synthesising (**1**) using ethanol. **B:**  $^{19}\text{F}\{^1\text{H}\}$  NMR of (**1**).

**A:**



**B:**



**Figure S2:** Crystal packing diagram of (**1**) highlighting the  $\pi\cdots\pi$  interactions.

