

Supplementary Material

Reactivity of β -diketiminato magnesium alkyl complexes: heterocumulenes and phosphanes

Morgan J. Taylor,^A Martyn P. Coles,^{A,B} and J. Robin Fulton^{A,B,C}

^ADepartment of Chemistry, University of Sussex, Falmer, Brighton, BN1 9QJ, UK.

^BCurrent address: School of Chemical and Physical Sciences, Victoria University of Wellington, Wellington 6012, NZ.

^CCorresponding author. Email: j.robin.fulton@vuw.ac.nz

Fig S1	^1H NMR spectrum of $[(\text{BDI})\text{Mg}(\text{MeC}\{\text{NCy}\}_2)]$ (3)	S2
Fig S2	^{13}C NMR spectrum of $[(\text{BDI})\text{Mg}(\text{MeC}\{\text{NCy}\}_2)]$ (3)	S3
Fig S3	Thermal ellipsoid plot of $[(\text{BDI})\text{Mg}(\text{MeC}\{\text{NCy}\}_2)]$ (3)	S4
Fig S4	^1H NMR spectrum of $[(\text{BDI})\text{Mg}(\text{BnC}\{\text{NCy}\}_2)]$ (4)	S5
Fig S5	^{13}C NMR spectrum of $[(\text{BDI})\text{Mg}(\text{BnC}\{\text{NCy}\}_2)]$ (4)	S6
Fig S6	^1H NMR spectrum of $[(\text{BDI})\text{Mg}(\text{PPh}_2)(\text{THF})]$ (5)	S7
Fig S7	^{13}C NMR spectrum of $[(\text{BDI})\text{Mg}(\text{PPh}_2)(\text{THF})]$ (5)	S8
Fig S8	Thermal ellipsoid plot of $[(\text{BDI})\text{Mg}(\text{PPh}_2)(\text{THF})]$ (5)	S9
Fig S9	^1H NMR spectrum of $[(\text{BDI})\text{Mg}(\text{PCy}_2)(\text{THF})]$ (6)	S10
Fig S10	^{13}C NMR spectrum of $[(\text{BDI})\text{Mg}(\text{PCy}_2)(\text{THF})]$ (6)	S11

Figure S1. ^1H NMR spectrum of $[(\text{BDI})\text{Mg}(\text{MeC}\{\text{NCy}\}_2)]$ (3)




Figure S2. ^{13}C NMR spectrum of [(BDI)Mg(MeC{NCy}₂)₂] (**3**).




Figure S3. Thermal ellipsoid plot of $[(\text{BDI})\text{Mg}(\text{MeC}\{\text{NCy}\}_2)]$ (**3**).




Figure S4. ^1H NMR spectrum of $[(\text{BDI})\text{Mg}(\text{BnC}\{\text{NCy}\}_2)]$ (**4**).




Figure S5. ^{13}C NMR spectrum of $[(\text{BDI})\text{Mg}(\text{BnC}\{\text{NCy}\}_2)]$ (**4**).




Figure S6. ^1H NMR spectrum of $[(\text{BDI})\text{Mg}(\text{PPh}_2)(\text{THF})]$ (**5**).




Figure S7. ^{13}C NMR spectrum of $[(\text{BDI})\text{Mg}(\text{PPh}_2)(\text{THF})]$ (**5**).




Figure S8. Thermal ellipsoid plot of $[(\text{BDI})\text{Mg}(\text{PPh}_2)(\text{THF})]$ (**5**).




Figure S9. ^1H NMR spectrum of [(BDI)Mg(PCy₂)(THF)] (**6**).




Figure S10. ^{13}C NMR spectrum of [(BDI)Mg(PCy₂)(THF)] (**6**).

