

SUPPLEMENTARY MATERIAL

Crystal Structure of a Heterometallic Luminophore – the Ru(II) Complex of a Ferrocenyl-terpyridine with a Flexible Linkage

Young Hoon Lee,^A Jong Won Shin,^B Yusuke Sekimoto,^C Shinya Hayami,^C Jack Harrowfield,^D Yang Kim^{E,F}

^A Department of Chemistry and EHSRC, University of Ulsan, Ulsan, 93 Daehak-ro, Nam-gu, Ulsan 44610, South Korea.

^B Daegu-Gyeongbuk Branch, Korea Institute of Science and Technology Information ,90 Yutongdanji-ro, Buk-gu, Daegu 41515, South Korea.

^C Department of Chemistry, Graduate School of Science and Technology, Kumamoto University, 2-39-1 Kurokami, Kumamoto 860-8555, Japan.

^D ISIS, Université de Strasbourg, 8 allée Gaspard Monge, 67083 Strasbourg, France.

^E Department of Chemistry & Advanced Materials, Kosin University, 194, Wachi-Ro, Yeongdo-gu, Busan 49104, South Korea.

^F Corresponding author. E-mail : ykim@kosin.ac.kr




Figure S1. ¹H NMR spectrum of compound 2 in CDCl₃ at 25 °C.




Figure S2. ¹H NMR spectrum of L1 in CDCl₃ at 25 °C.




Figure S3. ¹H NMR spectrum of [Ru(tpy)(L1)](ClO₄)₂ in CDCl₃ at 25 °C.