

**Supplementary material for**

**Prevalence and characteristics associated with concurrent smoking and alcohol misuse within Australian general practice patients**

*Breanne Hobden,<sup>1,2,3,4</sup> Jamie Bryant,<sup>1,2,3</sup> Kristy Forshaw,<sup>1,2,3</sup> Christopher Oldmeadow,<sup>3</sup> Tiffany-Jane Evans<sup>3</sup> and Rob Sanson-Fisher<sup>1,2,3</sup>*

<sup>1</sup>Health Behaviour Research Collaborative, HMRI Building, University of Newcastle, Callaghan, NSW 2308, Australia. Email: jamie.bryant@newcastle.edu.au; kristy.forshaw@newcastle.edu.au; Rob.Sanson-Fisher@newcastle.edu.au

<sup>2</sup>Priority Research Centre for Health Behaviour, HMRI Building, University of Newcastle, Callaghan, NSW 2308, Australia

<sup>3</sup>Hunter Medical Research Institute, HMRI Building, University of Newcastle, Callaghan, NSW 2308, Australia. Email: christopher.oldmeadow@hmri.org.au; tiffany.evans@hmri.org.au

<sup>4</sup>Corresponding author. Email: Bree.Hobden@newcastle.edu.au

**Table S1. Adjusted odds ratios for interaction of chronic condition with smoking – Random effects model.**

Variable	Category	OR (95% CI)	P-Value
Gender	Female vs Male	0.33 (0.3, 0.4)	<.0001
Age	30-44 vs 18-29	0.31 (0.2, 0.4)	<.0001
	45-64 vs 18-29	0.15 (0.1, 0.2)	
	65+ vs 18-29	0.04 (0, 0.1)	
SEIFA	Over 6 vs 6 or less	1.15 (0.9, 1.4)	0.1661
Education	TAFE/Dip vs High school and below	0.91 (0.7, 1.1)	0.0834
	University/Post Grad vs High school and below	0.8 (0.7, 1)	
	Other vs High school and below	0.79 (0.5, 1.3)	
ethnicity	ATSI vs Non Indigenous	1.09 (0.5, 2.5)	0.8393
Overweight or obese	Yes vs No	1.3 (1.1, 1.5)	0.0011
			<b>P-Value for interaction</b>
Chronic condition Yes vs No	Smokers	1.55 (1.0, 2.4)	0.0387
	Non-smokers	0.97 (0.8, 1.2)	