

## Supplementary Material for

### **Cost-effectiveness of a complex intervention in general practice to increase uptake of long-acting reversible contraceptives in Australia**

*Milena Lewandowska*<sup>1,10</sup> MPhil, Research Fellow

*Richard De Abreu Lourenco*<sup>1</sup> PhD, Associate Professor of Health Economics

*Marion Haas*<sup>1,2</sup> MPH, PhD, Professor of Health Economics

*Cathy J. Watson*<sup>3</sup> PhD, Research Fellow

*Kirsten I. Black*<sup>4</sup> PhD, Professor of Gynaecology

*Angela Taft*<sup>5</sup> PhD, Professor of Nursing and Midwifery, Principal Research Fellow

*Jayne Lucke*<sup>6</sup> PhD, Adjunct Professor of Psychology and Public Health

*Kevin McGeechan*<sup>1,2</sup> PhD, Associate Professor of Public Health

*Kathleen McNamee*<sup>7,8</sup> Dr, Adjunct Senior Lecturer

*Jeffrey F. Peipert*<sup>9</sup> MD, PhD, Professor and Chair of Obstetrics and Gynaecology

*Danielle Mazza*<sup>3</sup> MD, Professor of General Practice

<sup>1</sup>Centre for Health Economics Research and Evaluation, University of Technology Sydney, NSW, Australia. Email: Richard.DeAbreuLourenco@chere.uts.edu.au; Marion.Haas@chere.uts.edu.au

<sup>2</sup>School of Public Health, University of Sydney, NSW, Australia. Email: Kevin.McGeechan@sydney.edu.au

<sup>3</sup>Department of General Practice, Monash University, Melbourne, Vic., Australia. Email: Cathy.Watson@monash.edu; Danielle.Mazza@monash.edu

<sup>4</sup>Royal Prince Alfred Hospital, University of Sydney, NSW, Australia. Email: Kirsten.Black@sydney.edu.au

<sup>5</sup>Judith Lumley Centre, La Trobe University, Melbourne, Vic., Australia. Email: A.Taft@latrobe.edu.au

<sup>6</sup>School of Psychology and Public Health, La Trobe University, Melbourne, Vic., Australia. Email: J.Lucke@latrobe.edu.au

<sup>7</sup>Family Planning Victoria, Vic., Australia. Email: kmcnamee@fpv.org.au

<sup>8</sup>Obstetrics and Gynaecology, Monash Health, Monash University, Melbourne, Vic., Australia.

<sup>9</sup>Department of Obstetrics and Gynaecology, Indiana University School of Medicine, Indianapolis, IN, USA. Email: jpeipert@iu.edu

<sup>10</sup>Corresponding author. Email: Milena.Lewandowska@chere.uts.edu.au

**Table S1. Characteristics of participants in the ACCORd trial**


<b>Characteristics</b>	<b>Intervention group</b>	<b>Control group</b>	<b>Total</b>
Family physicians (N)	25	35	57
Gender			
Male	2 (8.0)	4 (12.5)	6
Female	23 (92.0)	28 (87.5)	51
Age group			
25-34	3 (12.0)	2 (6.3)	5
35-54	17 (68.0)	24 (75.0)	41
≥55	5 (20.0)	6 (18.8)	11
Inserts intrauterine devices			
No	22 (88.0)	27 (84.4)	47
Yes	3 (12.0)	5 (15.6)	8
Inserts implants			
No	7 (28.0)	10 (31.3)	17
Yes	18 (72.0)	22 (68.8)	40
Implants inserted each month, n			
1-4	3 (12.0)	3 (9.4)	6
5-9	1 (4.0)	4 (12.5)	5
≥10	21 (84.0)	25 (78.1)	46
Specific training in contraception			
No	19 (76.0)	24 (75.0)	43
Yes	6 (24.0)	8 (25.0)	14
Trained-insert intrauterine devices			
No	15 (60.0)	21 (65.6)	36
Yes	10 (40.0)	11 (34.4)	21
Women participants (N)	307	433	740
Age, y			
16-24	104 (33.9)	163 (37.6)	267
25-34	111 (36.2)	173 (40.0)	284
35-45	92 (30.0)	97 (22.4)	189
Parity			
0	207 (67.4)	313 (72.3)	520
1	24 (7.8)	32 (7.4)	56
2	53 (17.3)	71 (16.4)	124
≥3	23 (7.5)	17 (3.9)	40
Long-acting reversible contraceptives use at baseline			
No	266 (87.2)	379 (87.5)	645
Yes	39 (12.8)	54 (12.5)	93
Marital status			
Married/de facto	133 (43.5)	184 (42.5)	317
Single	173 (56.5)	249 (57.5)	422
Household income			
≤\$600 per week	75 (30.4)	126 (35.3)	201
>\$600 per week	172 (69.6)	231 (64.7)	403

Characteristics	Intervention group	Control group	Total
Education			
Completed <12 y	99 (32.2)	144 (33.3)	243
Completed ≥12 y	208 (67.8)	289 (66.7)	497
Previous unintended pregnancy			
No	249 (81.1)	363 (83.8)	612
Yes	58 (18.9)	70 (16.2)	128
Previous abortion			
No	267 (87.0)	390 (90.1)	657
Yes	40 (13.0)	43 (9.9)	83

Abbreviations: N=number; y=year

Source: Mazza et al 2019, Table 1

**Figure S1. Tornado plot of results of sensitivity analysis**



Abbreviations: LARC= long-acting reversible contraceptive; NMC= non-medical contraceptive; OC= oral contraceptive;

TOP=termination of pregnancy; UP=unplanned pregnancy

Notes: 1. Intervention group; 2. Usual care group