
Supplementary material

Molecular phylogenetic analysis of Australian arid-zone oniscidean isopods (Crustacea : *Haloniscus*) reveals strong regional endemism and new putative species

Michelle T. Guzik^{A,H}, Danielle N. Stringer^A, Nicholas P. Murphy^B, Steven J. B. Cooper^{A,C}, Stefano Taiti^{D,E}, Rachael A. King^{A,C}, William F. Humphreys^{F,G} and Andrew D. Austin^A

^AAustralian Centre for Evolutionary Biology and Biodiversity, School of Biological Sciences, The University of Adelaide, North Terrace, SA 5005, Australia.

^BDepartment of Ecology, Environment and Evolution, La Trobe University, Bundoora, Vic. 3086, Australia.

^CSouth Australian Museum, North Terrace, Adelaide, SA 5000, Australia.

^DIstituto per lo Studio degli Ecosistemi, Consiglio Nazionale delle Ricerche, Via Madonna del Piano 10, 50019 Sesto Fiorentino (Firenze), Italy.


^EMuseo di Storia Naturale dell'Università, Sezione di Zoologia, La Specola', Via Romana 17, 50125 Florence, Italy.

^FWestern Australian Museum, Welshpool DC, WA 6986, Australia.

^GSchool of Animal Biology, The University of Western Australia, Crawley, WA 6009, Australia.

^HCorresponding author. Email: michelle.guzik@adelaide.edu.au

Fig S1. For *18S* data, BI analyses (Supplementary Material 1) showed that the relationships among clades were consistent with *COI*.



0.008