

## Supplementary Material

### **Deep-sea movement patterns of the Patagonian toothfish *Dissostichus eleginoides* Smitt in the Southwest Atlantic**

*Brendon Lee<sup>A,B,\*</sup>, Frane Skeljo<sup>A</sup>, Haseeb S. Randhawa<sup>A,C,D,E</sup> and Alexander Arkhipkin<sup>A</sup>*

<sup>A</sup>Fisheries Department, Falkland Islands Government, Stanley, PO Box 598, FIQQ 1ZZ, Falkland Islands


<sup>B</sup>Department of Ichthyology and Fisheries Science, Rhodes University, PO Box 94, Grahamstown 6140, South Africa

<sup>C</sup>Faculty of Life and Environmental Sciences, University of Iceland, Sturlugata 7, 102 Reykjavik, Iceland

<sup>D</sup>South Atlantic Environmental Research Institute, Stanley Cottage, Stanley, FIQQ 1ZZ, Falkland Islands

<sup>E</sup>New Brunswick Museum, 277 Douglas Avenue, Saint John, NB, E2K 1E5, Canada

\*Correspondence to: Email: [blee@naturalresources.gov.fk](mailto:blee@naturalresources.gov.fk); [brendon82.lee@gmail.com](mailto:brendon82.lee@gmail.com)



**Fig. S1.** Spatial distribution of longline fishing effort (number of lines set per  $0.1 \times 0.1^\circ$  grid square;  $n = 2236$ ) during the study period (June 2016–October 2021) in the Falkland Islands Conservation Zones.