Ptilotis plumulus, and eggs301 Eggs of Meliphaga longi-Eggs of Anthochaera lunulata<sup>303</sup> Eggs a lewinii<sup>304</sup> Eggs of Myzantha obscura Anthochaera lewinii<sup>304</sup> Acanthagenys rufogularis<sup>305</sup> 2 Amadina castanotis<sup>306</sup> Estrelda Bichenovii, and eggs<sup>307</sup> Eggs of coronoides Eggs of Strepera . . . ?308 1 Cracticus destructor309 Eggs of Gymnorhina tibicen<sup>310</sup> Eggs of some unidentified species.

Gilbert left Western Australia on December 20, 1843, on the schooner Timbo, vessel of 123 tons, for Launceston, via Adelaide. In the Perth Gazette of December 23, it is noted that the Timbo sailed on December 20 for Launceston, but

no passengers are mentioned.

That concludes the review of the work done by Gilbert in Western Australia, and it remains only for the writer to express his sincere thanks to those numerous correspondents who have assisted him in filling an historical want by doing justice to the efforts of one who was indeed the pioneer ornithologist of Western Australia.

Part 1, p. 112, line 1, for 1939 read 1938. p. 127, line 29, for 153 read 157.

301. Gould described the eggs of this species in the Handbook, I, pp. 516-7. They were collected at York. See part 2, footnote 121. 302. = Meliornis novæ-hollandiæ, Yellow-winged Honeyeater. In the Handbook Gould describes its nidification. Gould must have obtained the information from Gilbert, as he records the bird under the name longivostris, the Western Australian form, yet he makes no acknowledgment to Gilbert.

303. A. carunculata, Red Wattle-bird.
304. A. chrysoptera, Brush Wattle-bird.
305. In the Handbook Gould records having received a pair from

306. Gould records (Handbook, I, p. 419) that "It has also been found, though very sparingly, at Swan River."
307. This is a puzzling item. Fringilla bichenovii was the name given by Vigors and Horsfield to the Double-banded Finch. It seems hardly possible that Gilbert could have used the name for the Redeared Firetail (Zonæginthus oculatus), yet that appears to be the only species to which the entry can refer. In the Handbook Gilbert's field-notes on the Red-eared Firetail are quoted, but there is no mention of the eggs.

308. S. versicolor.
309. C. torquatus. See part 1, foc
310. G. dorsalis, Western Magpie. See part 1, footnote 22.

New Genus.-The Short-billed Petrel, formerly called Pterodroma brevirostris, now known as P. lugens of Kuhl 1820, has the bill very much compressed on the sides. That character is shared by no other species of Pterodromine Petrel and it can be made a diagnostic character for a new subgenus Lugensa and be known as Pterodroma (Lugensa) lugens (Kuhl 1820). — Gregory Mathews, Sydney,