

Figure 1 shows the effective double polarization term I_2/I_0 for pure states 3S_1 , 3D_1 , and 3D_3 . The $J = 3$ curve is characteristically different from those for $J = 1$, having two maxima at 40° and 90° with an intervening zero at 63° .




Fig. 1.—Effective double polarization term ($m = 2$) for pure states 3S_1 , 3D_1 , and 3D_3 .

Acknowledgments

The author thanks Professor R. D. Ehrlich and Professor V. W. Hughes for discussion.

References

- BENVENUTI, A., CLINE, D., RUTZ, R., REEDER, D. D., and SCHERER, V. R. (1971).—*Phys. Rev. Lett.* **27**, 283.
 BURROWS, R. D., *et al.* (1970).—*Aust. J. Phys.* **23**, 819.
 CARSON, R. G., BUTTON-SCHAFER, J., HERTZBACH, S. S., KOFLER, R. R., and YAMAMOTO, S. S. (1972).—*Proc. 16th Int. Conf. on High Energy Physics, Chicago, Paper No. 498.*
 CLINE, D., and RUTZ, R. (1972).—*Phys. Rev. D* **5**, 778.
 PEASLEE, D. C. (1972).—*Particles Nuclei* **4**, 73.
 PEASLEE, D. C. (1973).—Straightness of boson trajectories. *Lett. Nuovo Cim.* (in press).

CORRIGENDUM

VOLUME 25, NUMBER 5

“Classification of supernova remnants and HII regions from their recombination line emission.” By J. R. Dickel and D. K. Milne. pp. 539–44

The galactic source number designations G35.6–0.4 and G35.5–0.0 in Table 1 should be interchanged. Thus G35.6–0.4 is the supernova remnant and G35.5–0.0 appears to be an HII region. The authors thank Dr. T Velusamy for calling this error to their attention.