

Supplementary Material

Resolution of the *Eremophila tietkensis* (Scrophulariaceae) species complex based on congruence between morphological and molecular pattern analyses

Amy L. Curtis^{A,B,*}, *Pauline F. Grierson*^B, *Jacqueline Batley*^B, *Jeremy Naaykens*^C, *Rachael M. Fowler*^D, *Anita Severn-Ellis*^B and *Kevin R. Thiele*^{A,B}

^AWestern Australian Herbarium, Biodiversity and Conservation Science, Department of Biodiversity, Conservation and Attractions, Locked Bag 104, Bentley Delivery Centre, WA 6983, Australia

^BSchool of Biological Sciences, Faculty of Science, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia

^CRio Tinto Iron Ore, 152–158 St Georges Terrace, Perth, WA 6000, Australia

^DSchool of BioSciences, The University of Melbourne, Parkville, Vic. 3010, Australia

*Correspondence to: Email: amy.curtis@dbca.wa.gov.au




Fig. S1. Quantity ($\text{ng } \mu\text{L}^{-1}$) and quality score of DNA extracted from herbarium specimens shown by year of collection. B) Sequence size (b) after demultiplexing and number of base pairs (kilobase pairs) of herbarium specimens shown by year of collection. Error bars represent standard deviation.