

Uptake of ‘Eradicat’ feral cat baits by non-target species on Kangaroo Island

Rosemary Hohnen^{A,E}, Brett P. Murphy^A, Sarah M. Legge^{B,C}, Chris R. Dickman^D and John C. Z. Woinarski^A

^ANational Environmental Science Program Threatened Species Recovery Hub, Research Institute for the Environment and Livelihoods, Ellengowan Drive, Charles Darwin University, Casurina, NT 0909, Australia.

^BNational Environmental Science Program Threatened Species Recovery Hub, Centre for Biodiversity and Conservation Science, Level 5 Goddard Building, University of Queensland, St Lucia, Qld 4072, Australia.

^CNational Environmental Science Program Threatened Species Recovery Hub, Fenner School, Building 141, Linnaeus Way, The Australian National University, Canberra, ACT 2601, Australia.

^DNational Environmental Science Program Threatened Species Recovery Hub, Desert Ecology Research Group, Heydon-Laurence Building, The University of Sydney, 325, A08, Camperdown, NSW 2006, Australia.

^ECorresponding author. Email: rosemary.hohnen@cdu.edu.au

Table S1. Equations used to calculate how much bait a species could use before reaching its LD50

Description	Equation
Amount of 1080 for LD50	$(\text{Bodyweight}) * (\text{LD50 mg kg}^{-1})$
Weight of bait for LD50	$((\text{Amount of 1080 for LD50}) / 4.5 \text{ mg amount of 1080 in bait}) * 15\text{g dry bait weight}$
Number of baits for LD50	$\text{Weight of bait for LD50} / 15\text{g dry bait weight}$