

[10.1071/AN21560](https://doi.org/10.1071/AN21560)

Animal Production Science

Supplementary Material

Can measurements of foraging behaviour predict variation in weight gains of free-ranging cattle?

David J. Augustine^{A}, Edward J. Raynor^A, Sean P. Kearney^A, and Justin D. Derner^B*

^AUnited States Department of Agriculture, Agricultural Research Service, Rangeland Resources and Systems Research Unit, Fort Collins, CO 80526, USA.

^BUnited States Department of Agriculture, Agricultural Research Service, Rangeland Resources and Systems Research Unit, Cheyenne, WY 82009, USA.

*Correspondence to: David J. Augustine United States Department of Agriculture, Agricultural Research Service, Rangeland Resources and Systems Research Unit, Fort Collins, CO 80526, USA Email: David.Augustine@usda.gov

Fig. S1. Yearling steers with MOOnitor collars grazing shortgrass rangeland in northeastern Colorado, USA. The steer on the left is with the collar centred correctly on the neck; on the right is an example of the solar panels shifted off-centre from the neck.



Fig. S2. Comparison of estimates of grazing hours per day calculated based on classification of 4 sec versus 5 min intervals for yearling steers grazing shortgrass rangeland in northeastern Colorado, USA.

