

Supplementary material

Reliability of leaf functional traits after delayed measurements

Dinesh Thakur^{A,B}, Lakhbeer Singh^A and Amit Chawla^{A,B,C}

^ACouncil of Scientific and Industrial Research (CSIR)–Institute of Himalayan Bioresource Technology (IHBT), Palampur, Kangra, Himachal Pradesh, 176061, India.

^BAcademy of Scientific and Innovative Research (AcSIR), CSIR–IHBT, Palampur, India.

^CCorresponding author. Email: amitchawla21@gmail.com




Fig. S1. Changes in leaf area (LA) with storage time for different species. D1, D2, etc. represent Day 1, Day 2, etc. Figure was prepared using 'ggline()' function in R.






Fig. S2. Changes in leaf rehydration percentage with storage time for different species. D1, D2, etc. represent Day 1, Day 2, etc. Figure was prepared using 'ggline()' function in R.




Fig. S3. Changes in specific leaf area (SLA) with storage time for different species. D1, D2, etc. represent Day 1, Day 2, etc. Figure was prepared using 'ggline()' function in R.





Fig. S4. Changes in leaf dry matter content (LDMC) with storage time for different species. D1, D2, etc. represent Day 1, Day 2, etc. Figure was prepared using 'ggline()' function in R.