

10.1071/CH11444\_AC

©CSIRO 2012

Australian Journal of Chemistry 2012, 65(3), 314–319.

## **Supplementary Material**


### **Highly Monodispersed PEG-stabilized Ni Nanoparticles: Proficient Catalyst for the Synthesis of Biologically Important Spiropyrans**

*Jitender M. Khurana<sup>A,B</sup> and Sneha Yadav<sup>A</sup>*


<sup>A</sup>Department of Chemistry, University of Delhi, Delhi 110007, India.

<sup>B</sup>Corresponding author. Email: jmkhurana1@yahoo.co.in


*2-amino-7-methyl-1',3',5-trioxo-1',3',5,6,7,8-hexahydrospiro[chromene-4,2'-indene]-3-carbonitrile* **4b**




**4b**




**4b**




*7'-amino-1',3'-dimethyl-1,2',3,4'-tetraoxo-1,1',2',3,3',4'-hexahydrospiro[indene-2,5'-pyrano[2,3-d]pyrimidine]-6'-carbonitrile 4e*




**4e**




*2-amino-1',3',5-trioxo-1',3',6,7-tetrahydro-5H-spiro[cyclopenta[b]pyran-4,2'-indene]-3-carbonitrile* **4g**




**4g**




*Ethyl 2'-amino-3'-cyano-6'-methyl-1,3-dioxo-1,3-dihydrospiro[indene-2,4'-pyran]-5'-carboxylate* **4l**




4l




4I




*2'-amino-7'-methyl-2,5'-dioxo-5',6',7',8'-tetrahydro-2H-spiro[acenaphthylene-1,4'-chromene]-3'-carbonitrile* **5b**




**5b**




**5b**




*2'-amino-7'-methyl-2,5'-dioxo-2H,5'H-spiro[acenaphthylene-1,4'-pyrano[4,3-b]pyran]-3'-carbonitrile* **5c**




**5c**




*7'-amino-1',3'-diethyl-2,4'-dioxo-2'-thioxo-1',2',3',4'-tetrahydro-2H  
spiro[acenaphthylene-1,5'-pyrano[2,3-d]pyrimidine]-6'-carbonitrile 5e*




**5e**




*2'-amino-2,5'-dioxo-6',7'-dihydro-2H,5H-spiro[acenaphthylene-1,4'-cyclopenta[b]pyran]-3'-carbonitrile* **5f**




**5f**



*2-amino-2',5,10-trioxo-5,10-dihydrospiro[benzo[g]chromene-4,3'-indoline]-3-carbonitrile* **6f**



**6f**

**6f**

