

Supplementary Materials

Selenium application influenced selenium biofortification and physiological traits in water-deficit common bean plants

Ruby Vega Ravello^{A,B}, Cynthia de Oliveira^A, Josimar Lessa^A, Lissa Vasconcellos Vilas Boas^C, Evaristo Mauro de Castro^C, Luiz Roberto Guimarães Guilherme^A and Guilherme Lopes^{A,D}

^ASoil Science Department, Federal University of Lavras UFLA, Lavras, Minas Gerais State, 37200-900, Brazil.

^BSoil Science Department, La Molina National Agrarian University – UNALM, La Molina 15024, Lima, Peru.

^CBiology Department, Federal University of Lavras, Lavras, Minas Gerais State, 37200-9000, Brazil.

^DCorresponding author. Email: guilherme.lopes@ufla.br




Fig. S1. Normalized transpiration ratio vs. fraction of transpirable soil water (FTSW) for non-irrigated treatments.




Fig. S2. N accumulation by grains (a), P accumulation by grains (b), K accumulation by grains (c), Mg accumulation by grains (d), and S accumulation by grains (e) of common bean concerning Se rates and soil water conditions. The vertical bars indicate standard errors of average values ($n = 4$). * and ***, Statistically significant difference at $p \leq 0.05$ and $p \leq 0.001$; ns, not significant.