

Australian Journal of Agricultural Research

Contents

Volume 58

Issue 9

2007

Supplement to 'Global Landscapes in Cereal Rust Control'

Leaf rust of spring wheat in Northern Kazakhstan and Siberia: incidence, virulence, and breeding for resistance.

A. Morgounov, L. Rosseeva, M. Koyshibaev 847

Sustained genetic control of wheat rust diseases in north-eastern Australia.

G. J. Platz, J. A. Sheppard 854

Crop/pasture improvement and protection

QTL analysis of malting quality traits in two barley populations.

J. F. Panozzo, P. J. Eckermann, D. E. Mather, D. B. Moody, C. K. Black, H. M. Collins, A. R. Barr, P. Lim, B. R. Cullis 858

Simple sequence repeat markers associated with three quantitative trait loci for black point resistance can be used to enrich selection populations in bread wheat.

M. J. Christopher, P. M. Williamson, M. Michalowicz, R. Jennings, A. Lehmensiek, J. Sheppard, P. Banks 867

Crop/pasture agronomy and physiology

Yield performance and adaptation of some Australian-grown rice varieties through multivariate analysis.

Siva Sivapalan, Graeme Batten, Ashantha Goonetilleke, Serge Kokot 874

Effect of second-pond dairy effluent on turnip dry matter yield, nutritive characteristics, and mineral content.

J. L. Jacobs, G. N. Ward 884

The effect of endophyte infection on persistence of tall fescue (*Festuca arundinacea* Schreb.) populations in two climatically contrasting Italian locations.

L. Pecetti, M. Romani, A. M. Carroni, P. Annicchiarico, E. Piano 893

Farming systems

A comparison of the establishment, productivity, and feed quality of four cocksfoot (*Dactylis glomerata* L.) and four brome (*Bromus* spp.) cultivars, under leaf stage based defoliation management.

L. R. Turner, D. J. Donaghy, P. A. Lane, R. P. Rawnsley 900

Grazing, mulching, and removal of wheat straw in a no-till system in a semi-arid environment.

S. Landau, Iris Schoenbaum, D. Barkai, E. D. Ungar, A. Genizi, J. Kigel 907

Animal nutrition and physiology

Responses of wool growth rate and body reserves to nutrition in Merino ewes: a potential biological link between production and fitness.

N. R. Adams, J. R. Briegel, J. C. Greeff 913

Animal breeding and genetics

Inclusion of skin follicle traits in selection indices in breeding programs improves genetic gain in Australian fine-wool Merinos.

M. Asadi Fozi, J. H. J. Van der Werf, A. A. Swan 921

Genetic evaluation of crossbred lamb production. 5. Age of puberty and lambing performance of yearling crossbred ewes.

N. M. Fogarty, V. M. Ingham, A. R. Gilmour, R. A. Afolayan, L. J. Cummins, J. E. Hocking Edwards, G. M. Gaunt 928