

[10.1071/FP22285](https://doi.org/10.1071/FP22285)

Functional Plant Biology

Supplementary Material



Exogenous *myo*-inositol increases salt tolerance and accelerates CAM induction in the early juvenile stage of the facultative halophyte *Mesembryanthemum crystallinum* but not in the late juvenile stage

Cheng-Hsun Li^A, Yun-Cheng Tu^A, Meng-Fang Wen^A, Hsing-Jung Tien^A, and Hungchen Emilie Yen^{A,}*

^ADepartment of Life Sciences, National Chung Hsing University, Taichung 40227, Taiwan.

*Correspondence to: Hungchen Emilie Yen Department of Life Sciences, National Chung Hsing University, Taichung 40227, Taiwan Email: heyen@dragon.nchu.edu.tw

Supplementary Fig. S1. Photosynthetic light-response curve (*A/I* curve) of early (*a*) and late (*b*) juvenile primary leaves.



Supplementary Fig. S2. Hydroponically-grown late juvenile plants with 5 pairs of expanded primary leaves at Day 0 (a) and early juvenile plants with 4 pairs of expanded primary leaves at Day 0 (b). The first pair of primary leaves was small, lanceolate and underneath the 3rd pair of primary leaves, therefore, was not visible in the pictures. The leaf pairs and growth stages were defined according to Adams *et al.* (1998). When the 7th leaf pair appears, growth of the primary axis terminates, the side shoots emerge, and the transition from juvenile to adult growth occurs.

(a)



(b)

