

Supplementary material

Preliminary evidence of spawning phenologies of freshwater fish in a wet–dry tropical river: the importance of both wet and dry seasons

A. J. King^{A,B}, C. Doidge^A, D. Buckle^A and K. J. Tyler^A

^AResearch Institute for Environment and Livelihoods, Charles Darwin University,
Darwin, NT 0909, Australia.

^BCorresponding author. Email: alison.king@cdu.edu.au

Table S1. Raw number and percentage of individuals across collection methods

TL, total larvae; J, juvenile; D, drift; LT, light trap; SNE, sweep-net electrofishing

		Raw numbers of individuals			Grand Total	Percentage		
		D	LT	SNE		D	LT	SNE
<i>Ambassis</i> spp.	TL	27	3	41	71	38	4	58
	J	64	186	38	288	22	65	13
<i>Amniataba percoides</i>	TL	58	63	9	130	45	48	7
	J	3	5	113	121	2	4	93
<i>Arramphus sclerolepsis</i>	J	0	0	5	5	0	0	100
<i>Craterocephalus</i> spp.	TL	8	233	325	566	1	41	57
	J	0	7	23	30	0	23	77
<i>Glossamia aprion</i>	TL	1	3	3	7	14	43	43
	J	1	9	14	24	4	38	58
<i>Glossogobius</i> spp.	J	0	0	18	18	0	0	100
<i>Hephaestus fuliginosus</i>	J	1	0	6	7	14	0	86
<i>Hypseleotris compressa</i>	TL	0	14	22	36	0	39	61
	J	0	8	57	65	0	12	88
<i>Leptachirus triramus</i>	J	1	0	13	14	7	0	93
<i>Leiopotherapon unicolor</i>	J	0	0	37	37	0	0	100
<i>Melanotaenia</i> spp.	TL	9	23	75	107	8	21	70
	J	3	21	76	100	3	21	76
<i>Mogurnda mogurnda</i>	J	0	0	3	3	0	0	100
<i>Nematalosa erebi</i>	TL	1	0	0	1	100	0	0
	J	8	0	9	17	47	0	53
<i>Neoarius graeffei</i>	J	0	0	4	4	0	0	100
<i>Neosilurus ater</i>	TL	1	0	0	1	100	0	0
	J	1	0	1	2	50	0	50
<i>Neosilurus hyrtlui</i>	J	5	0	7	12	42	0	58
<i>Ophisternon gutturale</i>	J	1	0	1	2	50	0	50
<i>Oxyeleotris lineolatus</i>	TL	6	0	0	6	100	0	0
	J	0	0	9	9	0	0	100
<i>Oxyeleotris selheimi</i>	J	0	0	6	6	0	0	100
<i>Strongylura krefftii</i>	J	0	0	19	19	0	0	100
<i>Syncomistes butleri</i>	TL	1	0	0	1	100	0	0
	J	4	0	1	5	80	0	20
<i>Toxotes chatareus</i>	TL	4	0	1	5	80	0	20
	J	9	7	234	250	4	3	94
UniID <i>Heramphidae</i> spp.	TL	1	0	1	2	50	0	50
UniID <i>Neosilurus</i> spp.	TL	27	0	2	29	93	0	7
UniID spB	TL	1	0	0	1	100	0	0
UniID spC	TL	12	3	34	49	24	6	69
UniID spD	TL	17	2	23	42	40	5	55
UniID spE	TL	1	0	23	24	4	0	96
UniID spF	TL	0	0	4	4	0	0	100
UniID spH	TL	52	0	0	52	100	0	0
UniID damaged	TL	39	34	41	114	34	30	36
Total number		367	621	1298	2286	16	27	57
Species richness		21	9	28	30	70	30	93




Fig. S1. $\text{log}(x+1)$ abundance of all larvae captured on each sampling trip in (a) drift nets, (b) light trap and (c) sweep-net electrofisher (SNE) samples.




Fig. S2. Non-metric multidimensional scaling (MDS) plots for larval abundance ($\log(x+1)$), showing (a) season and (b) method.