

Supplementary material

Determining the associated risk of beach-washed logs and their origin at Milman Island, a nesting ground for the endangered hawksbill turtle (*Eretmochelys imbricata*)

Caitlin E. Smith^{A,B}, *James Walker*^C, *Karen Donnan*^C, *Kimberly Riskas*^B
and *Christine A. Madden Hof*^{A,B,D}

^AWWF-Australia, Level 4, 340 Adelaide Street, Brisbane, Qld 4000, Australia.

^BThe University of the Sunshine Coast, 90 Sippy Downs Drive, Sippy Downs, Qld 4556, Australia.

^CNorthern Australia Quarantine Strategy, 114 Catalina Crescent, Cairns, Qld 4870, Australia.

^DCorresponding author. Email: chof@wwf.org.au




Figure S1. A heat map of hawksbill and green turtle nesting attempts as well as the distribution, length and circumference of logs around Milman Island. Vegetation types are shown around the island. Start points for each of the four sections are shown.

Table S1. The total abundance of logs and nesting attempts within four sections of nesting habitat on Milman Island, and their respective cumulative log length (m)

The ratio represents how many logs are present in each section per one nesting attempt

Section	Log abundance	Total nesting attempts	Ratio of logs to nesting attempts	Cumulative log length (m)
1	30	38	0.78 to 1	124.4
2	17	93	0.18 to 1	89.7
3	20	102	0.19 to 1	131
4	30	8	3.75 to 1	100.5