

## Corrigendum

Volume 36, Number 2

Early Life History of Barramundi, *Lates calcarifer* (Bloch), in North-eastern Queensland

*D. J. Russell and R. N. Garrett*

p. 197. Several names in the Prey item column of Table 2 required correction. For convenience, the complete table is reproduced below:

**Table 2. Gut contents of juvenile barramundi from nursery swamp and tidal creek habitats**  
Number of guts containing a particular food item and that item's frequency of occurrence in all stomachs that contained food are shown

Prey item	Nursery swamp (n = 85)		Tidal creek (n = 79)	
	No.	Frequency	No.	Frequency
<b>INSECTA</b>				
Chironomidae (larvae)	14	0.28	0	0
Culicidae (larvae)	4	0.08	0	0
Trichoptera (larvae)	1	0.02	0	0
Unidentified Insecta	4	0.08	0	0
<b>CRUSTACEA</b>				
Penaeidae	12	0.24	24	0.59
Portunidae	0	0	2	0.05
<b>OSTEICHTHYES</b>				
Hemirhamphidae				
<i>Zenarchopterus dispar</i>	1	0.02	0	0
Chandidae				
<i>Ambassis</i> sp.	1	0.02	13	0.32
Teraponidae				
<i>Amniataba caudavittatus</i>	2	0.04	0	0
<i>Terapon jarbua</i>	1	0.02	0	0
Scatophagidae				
<i>Selenotoca multifasciata</i>	0	0	1	0.02
Melanotaeniidae				
Melanotaeniidae sp.	0	0	1	0.02
Gobiidae				
Gobiidae sp.	9	0.18	4	0.10
Eleotridae				
Eleotridae sp.	1	0.02	1	0.02
<i>Hypseleotris</i> sp.	0	0	12	0.29
Siganidae				
<i>Siganus</i> sp.	1	0.02	0	0
Unidentified fish	8	0.16	19	0.46
Unidentified contents	2	0.04	0	0
No contents	35	—	38	—