

Supplementary Material

Adipocytokines may delay pubertal maturation of human Sertoli cells

I. V. Wagner^{A,B,D}, P. Yango^C, K. Svechnikov^A, N. D. Tran^C and O. Söder^A

^AKarolinska Institutet, Department of Women's and Children's Health, Pediatric Endocrinology Unit, 17176 Stockholm, Sweden.

^BDepartment of Pediatrics, University Hospital Cologne, Kerpener Str. 62, 50937, Cologne, Germany.

^CCenter for Reproductive Sciences, University of California, San Francisco, 513 Parnassus Avenue, CA 94143, USA.

^DCorresponding author. Email: isabel.wagner@ki.se

Table S1. Primer sequences

Name	Sequence	BP
<i>BMP4-F</i>	TGGGATTCCCGTCCAAGCTA	20
<i>BMP4-R</i>	ACGGAATGGCTCCATAGGTC	20
<i>CYP26A1-F</i>	CACCATTCTGGGATCTGGCT	20
<i>CYP26A1-R</i>	CCGCATAATCACCTTCTTGCG	21
<i>FGF-2-F</i>	CGGCTGTACTGCAAAAACGG	20
<i>FGF-2-R</i>	TTGTAGCTTGATGTGAGGGTCG	22
<i>FSHR-F</i>	CGC CCT CAT GGA AGC CAG CCT	21
<i>FSHR-R</i>	GCT CAG AGA TTT GCC GTC TCC AGT	24
<i>GAPDH-F</i>	TCC TGT TCG ACA GTC AGC CG	20
<i>GAPDH-R</i>	AGT TAA AAG CAG CCC TGG TGA	21
<i>GDNF-F</i>	GAACTCTTGCCCCTGACCTG	20
<i>GDNF-R</i>	GCGGCACCTCGGATCG	16
<i>LIF-F</i>	CCATACGCCACCCATGTCAC	20
<i>LIF-R</i>	CCCTGGGCTGTGTAATAGAGAA	22