

[10.1071/SR21013](https://doi.org/10.1071/SR21013)

Soil Research

Supplementary Material

Storage of soil samples leads to over-representation of the contribution of nitrate to plant-available nitrogen

Taleta Bailey^{A,B,}, Nicole Robinson^A, Mark Farrell^B, Ben Macdonald^C, Tim Weaver^D, Diogenes L. Antille^C, Aidan Chin^A, and Richard Brackin^A*

^ASchool of Agriculture and Food Sciences, The University of Queensland, Brisbane, Qld, Australia.

^BCSIRO Agriculture and Food, Waite Campus, Adelaide, SA, Australia.

^CCSIRO Agriculture and Food, Black Mountain, Canberra, ACT, Australia.

^DCSIRO Agriculture and Food, Myall Vale, Narrabri, NSW, Australia.

*Correspondence to: Taleta Bailey School of Agriculture and Food Sciences, The University of Queensland, Brisbane, Qld, Australia Email: taleta.bailey@uq.net.au

Supplementary material

Table S1. Soil temperature in each field plot of a cotton field trial (near Toobeah, Queensland, Australia) measured at the time of in situ microdialysis sampling (January 2019, 99 days after planting).

Field plot	Nitrogen rate (kg N ha⁻¹)	Temperature (°C)
3	172	24.1
4	172	24.2
5	309	24.3
6	309	24.2
7	172	25.0
8	172	26.0
9	309	26.0
10	309	26.1
15	172	26.0
16	172	26.1
17	309	26.2
18	309	26.4
20	309	26.8
21	309	27.0
22	172	27.0
23	172	27.0