

Supplementary Material

Implementation constraints limit benefits of restoration treatments in mixed-conifer forests

Jamie M. Lydersen^{A,D,E}, Brandon M. Collins^B and Carolyn T. Hunsaker^C

^AEcosystem Sciences Division, Department of Environmental Science, Policy, and Management, University of California, Berkeley, CA 94720, USA.

^BCenter for Fire Research and Outreach, University of California, Berkeley, CA 94720-3114, USA.

^CUSDA Forest Service, Pacific Southwest Research Station, Fresno, CA 93710, USA.

^DPresent address: California Department of Forestry and Fire Protection, Fire and Resource Assessment Program, Sacramento, CA 95818, USA.

^ECorresponding author. Email: jmlydersen@berkeley.edu

Table S1. Mortality models tests of fixed effects

BA, basal area

Effect	Proportion BA dead		Dead BA		Density dead trees	
	FValue	ProbF	FValue	ProbF	FValue	ProbF
Site	0.37	0.5858	0.01	0.9126	0.27	0.6047
Treatment	0.23	0.8724	0.30	0.8280	0.35	0.7922
Type	1.51	0.2216	0.61	0.4357	0.31	0.5777
Type × treatment	1.02	0.3881	0.79	0.5030	0.19	0.9038
Elevation	0.09	0.7603	0.32	0.5709	0.00	0.9668
Aspect	0.67	0.4141	0.20	0.6549	0.84	0.3602
Slope	1.72	0.1921	0.85	0.3586	0.63	0.4290
Post-thin live tree density	1.40	0.2397	0.04	0.8421	51.51	<0.0001