

Climatic correlates of migrant Natal long-fingered bat (*Miniopterus natalensis*) phenology in north-eastern South Africa

Mariëtte Pretorius^A, Hugh Broders^B, Ernest Seamark^C and Mark Keith^{A,D}

^AMammal Research Institute, Faculty of Natural and Agricultural Sciences, University of Pretoria, Hatfield 0028, Private Bag x20, South Africa.

^BDepartment of Biology, University of Waterloo, 200 University Avenue West, Waterloo, Ontario, N2 L 3G1, Canada.

^CAfricanBats NPC, 357 Botha Avenue, Kloofsig, 0157, South Africa.

^DCorresponding author. Email: mark.keith@up.ac.za

Table S1. Summary of total cumulative rainfall (Σ Rain) and mean average temperature (\bar{x} Temp) \pm standard error (SE) across the four year samples and four seasons recorded from the weather station at the Madimatle Cave, Limpopo, South Africa. Means in the last column followed by the same letter in bold are not significantly different ($p < 0.05$) according to Kruskal-Wallis pairwise post-hoc test

Year sample	Autumn		Winter		Spring		Summer		Total Rainfall	Pooled \bar{x} Temp
	Σ Rain	\bar{x} Temp \pm SE	Σ Rain	\bar{x} Temp \pm SE	Σ Rain	\bar{x} Temp \pm SE	Σ Rain	\bar{x} Temp \pm SE		
2014-2015	22.20	21.02 \pm 0.42	0	15.00 \pm 0.34	93.60	23.85 \pm 0.32	264.20	25.22 \pm 0.22	380.00	21.26 a
2015-2016	133.80	20.96 \pm 0.34	0.60	16.80 \pm 0.39	115.60	25.63 \pm 0.43	167.20	27.75 \pm 0.27	417.20	22.77 b
2016-2017	127.40	20.69 \pm 0.33	15.00	15.70 \pm 0.31	145.80	25.24 \pm 0.34	458.20	24.62 \pm 0.24	746.40	21.56 ac
2017-2018	77.60	20.83 \pm 0.37	2.80	16.20 \pm 0.23	131.20	24.02 \pm 0.32	302.80	25.24 \pm 0.23	514.40	21.57 abc
Pooled total/ mean	361.00	20.87	18.40	15.90	486.20	24.69	1192.40	25.71		