

[10.1071/WR22103](https://doi.org/10.1071/WR22103)

*Wildlife Research*

### **Supplementary Material**


#### **Detections of house mice on Gough Island approach zero within days of aerial baiting**

*Araceli Samaniego<sup>A,\*</sup>, Kim L. Stevens<sup>B</sup>, Vonica Perold<sup>B</sup>, Steffen Oppel<sup>B</sup>, and Pete McClelland<sup>B</sup>*

<sup>A</sup>Manaaki Whenua – Landcare Research, 231 Morrin Road, St Johns, Auckland 1072, New Zealand.

<sup>B</sup>RSPB Centre for Conservation Science, Royal Society for the Protection of Birds, The David Attenborough Building, Pembroke Street, Cambridge CB2 3QZ, UK.


\*Correspondence to: Araceli Samaniego Manaaki Whenua – Landcare Research, 231 Morrin Road, St Johns, Auckland 1072, New Zealand Email: [SamaniegoA@landcareresearch.co.nz](mailto:SamaniegoA@landcareresearch.co.nz)



## Gough Island Bait Supplementary Baiting

Scale at A4: 11:70,000  
Gough Island Restoration Program  
9/08/2021

Supplementary Material Figure S1. Bait application rates and area covered during the supplementary application on Gough Island in August 2021.



Supplementary Material Figure S2. Camera monitoring on Gough Island in 2021. 1 = last mouse detection at Base (19 June); 2 = last mouse detection during mouse monitoring (10 July); 3 = first mouse detection post-baiting operation (11 December).