

Supplementary material

Post-anthesis heat and a *Gpc-B1* introgression have similar but non-additive effects in bread wheat

Lancelot Maphosa^{A,B,C}, *Nicholas C. Collins*^{A,B}, *Julian Taylor*^B and *Diane E. Mather*^{A,B,D}

^AAustralian Centre for Plant Functional Genomics, Waite Research Institute, The University of Adelaide, PMB 1, Glen Osmond, SA 5064, Australia.

^BSchool of Agriculture, Food and Wine, Waite Research Institute, The University of Adelaide, PMB 1, Glen Osmond, SA 5064, Australia.

^CPresent address: Department of Environment and Primary Industries, 110 Natimuk Road, Horsham, Vic. 3400, Australia.

^DCorresponding author. Email: diane.mather@adelaide.edu.au




Fig. S1. Consensus-map positions (horizontal lines) of 3772 SNP markers that had previously been mapped (Cavanagh *et al.* 2013) on wheat chromosomes other than 6B and that were polymorphic between RAC1261A and Somerset and/or between Wyalkatchem and Burnside, indicating the positions at which polymorphisms were detected between RAC1262A and RS4-11-10 (S, indicating an allele derived from Somerset) and/or between Wyalkatchem and WB4-1-6 (B, indicating an allele derived from Burnside). The scale on the left indicates genetic distances in cM.