

10.1071/FP16036_AC

© CSIRO 2017

Supplementary Material: *Functional Plant Biology*, 2017, 44(9), 867–876.

Supplementary Material

Contrasting oxygen dynamics in *Limonium narbonense* and *Sarcocornia fruticosa* during partial and complete submergence

Elisa Pellegrini^{A,B,C}, *Dennis Konnerup*^B, *Anders Winkel*^B, *Valentino Casolo*^A and *Ole Pedersen*^B

^APlant Biology Unit, Department of Agricultural, Food, Environmental and Animal Science, University of Udine, via delle Scienze 206, 33100 Udine, Italy.

^BFreshwater Biological Laboratory, Department of Biology, University of Copenhagen, Universitetsparken 4, 2100 Copenhagen, Denmark.

^CCorresponding author. Email: elisa.pellegrini87@gmail.com


Table S1. Steady state intraplant O₂ concentrations (μmol L⁻¹) in photosynthetic tissues and roots in *Limonium narbonense* and *Sarcocornia fruticosa*, during drained condition, waterlogging, partial or complete submergence, in darkness or light

Values are means ± standard deviation ($n = 3$); ♦ indicates $n = 2$. Letters indicate significant differences ($P \leq 0.05$) based upon ANOVA and Tukey's HSD post-hoc test


	Darkness				Light			
	Drained condition	Waterlogging	Partial submergence	Complete submergence	Drained condition	Waterlogging	Partial submergence	Complete submergence
<i>L. narbonense</i>								
Petiole	172.4±4.9 ^a	167±25.2 ^a	9.1±6.1 ^g	4.5±4.9♦ ^g	174.2±4.7 ^a	163.1±3.1 ^{ab}	91.8±10.7 ^{cd}	73±28 ^{cdef}
Adventitious root	69±22.9 ^{cdef}	71.6±27 ^{cdef}	5.6±4.2 ^g	0±0 ^g	71.3±23.3 ^{cdef}	71.8±21.4 ^{cdef}	17.9±14.1 ^{fg}	6.6±4.5 ^g
<i>S. fruticosa</i>								
Article	194.6±18.4 ^a	198.4±25.3 ^a	106.8±28.1 ^{bc}	86.8±30.3 ^{cde}	195.6±18.6 ^a	204.7±23.3 ^a	180.1±7.4 ^a	179.7±13.7 ^a
Root	71.7±17 ^{cdef}	98.8±30.6 ^{cd}	44.4±16 ^{defg}	3.8±5 ^g	75.8±14.4 ^{cdef}	57.9±16.1 ^{cdefg}	31.3±3♦ ^{efg}	25.1±2 ^{fg}

muddy sediment (BB site)

Low stress (drained-waterlogged)




High stress (waterlogged-submerged)



sandy sediment (BO site)

Low stress (drained-waterlogged)



High stress (waterlogged-submerged)




Fig. S1. Oxygen dynamics, expressed as O₂ partial pressure, in sediment monitored during plant acclimation, in muddy sediment (BB) and sandy sediment samples (BO). In each graph, continuous vertical lines refer to an increase of water level (up to waterlogging for the low stress, up to complete submergence for high stress), dashed vertical lines refer to a decrease of water level (down to drained condition for low stress, down to waterlogging for high stress). The two species are identified by the abbreviations Lim for *Limonium narbonense* and Sar for *Sarcocornia fruticosa* with a number (1–6) corresponding to different plant individuals.