

Supplementary Material

No escape? Costs and benefits of leaf de-submergence in the pasture grass *Chloris gayana* under different flooding regimes

Gustavo G. Striker^{A,B,E,F}, Cecilia Casas^C, Xiaolin Kuang^{A,B} and Agustín A. Grimoldi^{A,D}

^AIFEVA, Universidad de Buenos Aires, CONICET, Facultad de Agronomía, Avenida San Martín 4453, CPA 1417, DSE Buenos Aires, Argentina.

^BCátedra de Fisiología Vegetal, Facultad de Agronomía, Universidad de Buenos Aires. Av. San Martín 4453 (CPA 1417 DSE) Buenos Aires, Argentina.

^CCátedra de Edafología, Facultad de Agronomía, Universidad de Buenos Aires. Av. San Martín 4453 (CPA 1417 DSE) Buenos Aires, Argentina.

^DCátedra de Forrajicultura, Facultad de Agronomía, Universidad de Buenos Aires. Av. San Martín 4453 (CPA 1417 DSE) Buenos Aires, Argentina.

^ESchool of Agriculture and Environment, Faculty of Science, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia.

^FCorresponding author. Email: striker@agro.uba.ar




Fig. S1. Picture showing the leaf-elongation ‘escape’ strategy developed by 2-week-old plants of *Chloris gayana* cultivars from Experiment 1. Photo was taken 3.5 days after submergence started. All cultivars promoted leaf-elongation showing leaves above water after being completely submerged at the beginning of the treatment.