

## Supplementary Material

### Uncovering early response to drought by proteomic, physiological and biochemical changes in the almond × peach rootstock ‘Garnem’

Beatriz Bielsa<sup>A</sup>, María Á. Sanz<sup>B</sup> and María J. Rubio-Cabetas<sup>A,C</sup>


<sup>A</sup>Unidad de Hortofruticultura, Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA) – IA2 (CITA-Universidad de Zaragoza), Av. Montañana 930, 50059, Zaragoza, Spain.

<sup>B</sup>Área de Laboratorios de Análisis y Asistencia Tecnológica, Centro de Investigación y Tecnología Agroalimentaria de Aragón (CITA), Av. Montañana 930, 50059, Zaragoza, Spain.

<sup>C</sup>Corresponding author. Email: [mjrubioc@cita-aragon.es](mailto:mjrubioc@cita-aragon.es)

File S1. Accumulation patterns throughout drought stress experiment for each differential spot identified by LC-EMI-MS/MS. Control plants: Dark grey bars and black solid lines; Stressed Plants: Light grey bars and black dash lines. (H = hour; R = Recovery).

**Figure S1.** Accumulation patterns throughout drought stress experiment for each differential spot identified by LC-EMI-MS/MS. Control plants: Dark grey bars and black solid lines; Stressed Plants: Light grey bars and black dash lines. (H = hour; R = Recovery).



**Differential spots at both 2 h and 24 h of drought stress**

