

Supplementary Material

Transcriptional responses of *Hypericum perforatum* cells to *Agrobacterium tumefaciens* and differential gene expression in dark glands

Weina Hou^{A,B}, *Rupesh K. Singh*^C, *Viviana Martins*^B, *Francisco Tenllado*^{D,F}, *Gregory Franklin*^A and *Alberto C. P. Dias*^{A,B,E,F}

^ACentre for the Research and Technology of Agro-Environmental and Biological Sciences (CITAB), Department of Biology, University of Minho, 4710-057, Braga, Portugal.

^BCentre of Molecular and Environmental Biology (CBMA), Department of Biology, University of Minho, 4710-057, Braga, Portugal.

^CCentro de Química de Vila Real (CQ-VR), UTAD, 5000-801, Vila Real, Portugal.

^DDepartamento de Biotecnología Microbiana y de Plantas, Centro de Investigaciones Biológicas Margarita Salas, CSIC, Madrid, 28040, Spain.

^ECenter of Biological Engineering (CEB), University of Minho, 4710-057, Braga, Portugal.

^FCorresponding authors. Emails: tenllado@cib.csic.es; acpdias@bio.uminho.pt




Fig. S1. Expression variation of genes involved in RNA silencing pathway in *H. perforatum* L. leaves with (HpWith) or without (HpWithout) glands. (a) Relative transcript level of *RDR6*, (b) *SGS3* and (c) *AGO5*, as detected by qRT-PCR. Asterisks denote the significance levels compared with control: ns denotes not significant.