

Supplementary Material

Copper and iron metal resistant rhizospheric bacteria boost the plant growth and bacoside A content in *Bacopa monnieri* under stress conditions

Ramesh Kumar Kushwaha^{A,*}, Samyukta Madhav Joshi^A, Renuka Bajaj^A, Anthati Mastan^B, Vinay Kumar^C, Himani Patel^D, S. Jayashree^A, and Satya Prakash Chaudhary^E

^ADepartment of Biochemistry, School of Allied Health Sciences, REVA University, Bangalore 560064, Karnataka, India.

^BMicrobial Technology Laboratory, CSIR (Council of Scientific and Industrial Research)-Central Institute of Medicinal and Aromatic Plants, Research Center, Bangalore 560065, Karnataka, India.

^CPlant Genetic Resources and Improvement Division, CSIR-National Botanical Research Institute, Lucknow 226001, Uttar Pradesh, India.


^DBirbal Sahni Institute of Palaeosciences, 53 University Road, Lucknow 226007, Uttar Pradesh, India.

^EDepartment of Dravyagun, IMS (Institute of Medical Sciences), Banaras Hindu University, Varanasi 221005, Uttar Pradesh, India.


*Correspondence to: Ramesh Kumar Kushwaha Department of Biochemistry, School of Allied Health Sciences, REVA University, Bangalore 560064, Karnataka, India Email: Kushwaha.rameshkumar@gmail.com

Supplementary Figure S1.


(a) Chromatogram of standard bacoside A




(b) Chromatogram of control plants




(c) Chromatogram of Cu-1 treated plants




(d) Chromatogram of Cu-2 treated plants




(e) Chromatogram of Cu-3 treated plants




(f) Chromatogram of Cu-4 treated plants



(g) Chromatogram of Fe-1 treated plants



(h) Chromatogram of Fe-2 treated plants



(i) Chromatogram of Fe-3 treated plants

