

[10.1071/FP23312](https://doi.org/10.1071/FP23312)

Functional Plant Biology

Supplementary Material

Exogenous *Streptomyces* spp. enhance the drought resistance of naked oat (*Avena nuda*) seedlings by augmenting both the osmoregulation mechanisms and antioxidant capacities

Meixia Qiao^A, Siyuan Lv^A, Yuejing Qiao^{A,B}, Wen Lin^{A,B}, Zhiqiang Gao^{A,B}, Xiwang Tang^C, Zhenping Yang^{A,B,}, and Jie Chen^{A,B,*}*

^ACollege of Agriculture, Shanxi Agricultural University, Taigu, Shanxi 030801, China.

^BMinisterial and Provincial Co-Innovation Centre for Endemic Crops Production with High-quality and Efficiency in Loess Plateau, Taigu, Shanxi 030801, China.

^CHebei Key Laboratory of Agroecological Safety, Hebei University of Environmental Engineering, Qinhuangdao, Hebei 066102, China.

*Correspondence to: Zhenping Yang College of Agriculture, Shanxi Agricultural University, Ministerial and Provincial Co-Innovation Centre for Endemic Crops Production with High-quality and Efficiency in Loess Plateau, Taigu, Shanxi, China Email: yangzhenping340@163.com Jie Chen College of Agriculture, Shanxi Agricultural University, Ministerial and Provincial Co-Innovation Centre for Endemic Crops Production with High-quality and Efficiency in Loess Plateau, Taigu, Shanxi, China Email: chenjie03306@163.com

Supplementary materials:

Supplementary Table S1. Principal component analysis of each index under different treatments of S5 condition

Indexes	Principal Component					
	PC1	PC2	PC3	PC4	PC5	PC6
Shoot dry weight	0.388	-0.676	0.456	0.314	-0.103	0.253
Root dry weight	0.477	-0.501	-0.384	0.560	0.182	0.147
Root/Shoot	-0.182	0.758	-0.110	0.448	0.399	0.097
Shoot moisture content	-0.029	0.228	0.855	-0.357	0.114	0.214
Root moisture content	-0.058	0.430	-0.483	0.573	0.111	0.473
Root length	-0.945	-0.234	0.168	-0.082	0.041	-0.035
Root surface area	-0.887	-0.112	0.192	0.019	0.184	0.322
Root volume	-0.627	0.180	0.242	-0.203	0.636	0.125
Root average diameter	0.493	0.168	-0.434	-0.705	-0.108	0.154
Root tips	0.281	0.841	0.089	-0.324	0.300	0.028
Root forks	-0.917	-0.340	-0.031	0.000	-0.122	0.037
Root cross	-0.924	-0.291	0.002	0.061	-0.112	0.111
TTC	0.700	-0.601	0.058	-0.180	0.285	-0.005
Root Pro	0.902	-0.146	-0.281	0.044	-0.110	-0.155
Root soluble sugar	0.700	-0.081	0.019	0.243	0.653	0.006
Root CAT	0.457	-0.283	-0.373	-0.645	0.319	0.148
Root POD	0.577	-0.400	0.343	-0.105	-0.297	0.532
Leaf Pro	0.412	0.620	-0.110	-0.061	-0.497	0.427
Leaf soluble sugar	0.684	0.039	0.401	-0.009	0.252	0.038
Leaf CAT	0.710	0.020	0.574	0.347	0.060	-0.148
Leaf POD	0.367	0.645	0.442	0.218	-0.417	-0.176
Characteristic Value	8.118	3.986	2.691	2.415	1.994	1.090
Variance %	38.657	18.981	12.814	11.499	9.496	5.190
Accumulation %	38.657	57.638	70.452	81.951	91.447	96.637

Supplementary Table S2. Principal component analysis of each index under different treatments of S15 condition

Indexes	Principal Component				
	PC1	PC2	PC3	PC4	PC5
Shoot dry weight	0.940	-0.231	-0.038	0.106	0.105
Root dry weight	0.917	0.074	0.306	0.123	0.171
Root shoot ratio	-0.538	-0.013	0.595	-0.191	0.058
Shoot moisture content	-0.522	-0.156	-0.650	0.415	-0.316
Root moisture content	0.442	0.450	0.725	-0.027	0.086
Root length	0.512	0.751	-0.221	0.137	-0.086
Root surface area	0.747	0.582	-0.030	0.216	-0.071
Root volume	0.815	0.422	0.021	0.352	0.020
Root average diameter	-0.218	-0.837	0.358	0.138	0.137
Root tips	0.673	0.684	-0.021	-0.073	-0.101
Root forks	-0.192	0.767	0.151	0.389	-0.090
Root cross	-0.264	0.795	-0.320	0.044	0.408
TTC	0.706	-0.520	-0.395	0.082	0.104
Root Pro	0.887	-0.114	0.262	-0.109	-0.321
Root soluble sugar	0.756	0.017	0.029	-0.608	0.159
Root CAT	0.732	-0.134	-0.416	-0.384	0.149
Root POD	0.780	-0.138	-0.393	-0.312	0.209
Leaf Pro	0.820	-0.326	-0.024	-0.031	-0.439
Leaf soluble sugar	0.928	-0.220	0.132	0.045	-0.169
Leaf CAT	0.587	-0.512	0.171	0.518	-0.031
Leaf POD	0.408	-0.380	0.005	0.644	0.481
Characteristic Value	9.619	4.638	2.302	1.923	1.025
Variance %	45.806	22.085	10.960	9.158	4.882
Accumulation %	45.806	67.891	78.851	88.010	92.892